At the Edge of the Universe:
Latest results from the deepest astronomical surveys, 9-13 October 2006, Sintra (Portugal)
*ASP Conference Series, Vol. ***, 2006*
J. Afonso, H. Ferguson, R. Norris

The Statistical Properties Of The Very Weak Radio Source Population In The GOODS/ACS HDF-N Region

T.W.B. Muxlow

MERLIN-VLBI National Facility, Jodrell Bank Observatory, University of Manchester, United Kingdom

R.J. Beswick, H. Thrall, A.M.S Richards, S.T. Garrington, A. Pedlar

Jodrell Bank Observatory, University of Manchester, United Kingdom

arXiv:astro-ph/0612075v1 4 Dec 2006

Abstract. Deep combination radio observations at 1.4GHz with the VLA and MERLIN have imaged a region 10 arcminutes square surrounding the Hubble Deep Field North (HDF-N). Initial studies of the weak radio source population have shown that the proportion of starburst systems increases with decreasing radio flux density with more than 70% of radio sources being starburst in nature at flux densities less than $S_{1.4GHz}\sim70\mu$Jy. The recently published GOODS ACS field overlaps this area, and here we present the results of a follow-up statistical study of the very weak radio sources ($S_{1.4GHz}<40\mu$Jy) in an 8.5 arcminute square field centred on the HDF-N which contains the region of overlap. Radio emission at the level of a few μJy are statistically detected associated with ACS galaxies brighter than a z-band magnitude of 25. These very faint radio sources are extended starburst systems with average radii in the range 0.6 to 0.8 arcseconds and for those with measured redshifts, radio luminosities typically several times that of the nearby well-studies starburst galaxy M82.

1. Introduction

Deep galaxy studies at visible and infra-red wavelengths have indicated that early galaxies merge to form larger systems in a 'bottom-up' scenario of galaxy assembly. This implies that galaxy-galaxy interactions were common at early epochs. Such interactions at modest redshifts are seen to trigger major star-formation activity and it is well established that star-formation rate density increases dramatically out to redshifts of $\sim$1 (Madau, Pozzetti, & Dickinson 1998). The star-formation rate for such galaxies can be estimated from both the centimetric radio and far infra-red (FIR) luminosities which have been found to be tightly correlated over several orders of magnitude (van der Kruit 1973; Condon *et al.* 1982).

Differential radio source counts at 1.4GHz (See Figure 1) show an increase at flux densities below 1mJy (Seymour, McHardy, & Gunn 2004). Multi-wavelength studies of a number of fields including deep radio observations involving the VLA, ATCA, MERLIN, and the EVN, have shown that this new weak radio source population is associated with distant star-forming galaxies.



Figure 1. Normalised differential radio source counts at 1.4GHz. At flux densities below 1mJy there is an upturn which is identified with a new population of distant star-forming galaxies. After Seymour, McHardy & Gunn (2004).

2. The Initial Radio Study of the HDF-N

Deep radio observations of the HDF-N region were made with the VLA and MERLIN during 1996 and 1997. The VLA results were presented in Richards *etal*. (1998) and Richards (2000), and the combination high angular resolution results were presented in Muxlow *et al*. (2005). Within a 10 arcminute square field centred on the HDF, 92 radio sources were detected by the VLA at 1.4GHz above a completeness limit of 40μJy/beam (5.3σ) using a 2 arcsecond beam. Combination high-resolution MERLIN+VLA images involving 42 hours of VLA observations and 18 days of MERLIN data were made for small regions around each of the 92 detected sources. The rms noise level in the combination images was $\sim$3.3μJy per 0.2 arcsecond beam making these images some of the most sensitive yet produced at 1.4GHz.

The full results of the study of the individual radio sources in the 10 arcminute field surrounding the HDF-N are given in Muxlow *et al*. (2005) and are summarized below:

• Within the 10x10 arcminute field there are 92 radio sources with flux densities at 1.4GHz >40μJy.

• The radio sources have angular sizes in the range 0.2 to 3 arcseconds.

• 85% of the radio sources are associated with galaxies brighter than 25$^{\rm th}$ magnitude in the v-band.

• The remaining 15% are optically faint or extremely red objects lying at high redshift, some of which are detected at sub-mm wavebands.

• The radio sources can be characterised as AGN or starburst according to their radio structures with starburst systems being identified as those with steep radio spectral indices and radio structures extended on (sub-)galactic scales overlaying the central region of the optical galaxy.
• Below 60μJy the radio sources are dominated by starburst type systems (See Figure 2).
• The starburst systems typically have radio luminosities significantly greater than that of nearby well-studied starbursts such as M82 or Arp220.
• Some high redshift starburst systems show evidence for embedded AGN (Garrett *et al.* 2001).



Figure 2. Distribution of source classification with flux density for the radio sources in the 10 arcminute field. After Muxlow *et al.*(2005).

3. A New Study of the Very Weak Radio Source Population

The initial investigation has been extended to include a statistical study of the very weak radio source population in the HDF-N region utilising ancillary data obtained from the large multi-wavelength Great Observatories Origins Survey (GOODS). The GOODS North region partially overlies the original field, and a new wide-field contiguous combination radio image was made covering a complete 8.5 arcminute square field centred on the HDF-N which contains the region of overlap. The 8.5 arcminute square radio image covers a region of the GOODS-N field containing 13030 galaxies brighter than magnitude 28.3 in the z-band ACS catalogue.

3.1. Faint Radio Emission from Optically-selected Galaxies

The radio and optical fields are astrometrically aligned to better than 50mas. Thus the radio flux density from each of the 13030 z-band optical galaxies can be investigated by integrating the radio emission from the MERLIN+VLA image of the 8.5 arcminute square field within a radius of 0.75 arcseconds at the position of every galaxy. Radio sources brighter than 40μJy have already been reported in detail in Muxlow *et al.* (2005) and were excluded from the sample. In addition, all galaxies with nearest neighbours closer than 1.5 arcseconds were also excluded so as to avoid confusion problems. The median flux densities, binned by z-band

magnitude are shown in Figure 3. A control sample was also constructed which incorporated a random 7 arcseconds positional shift between the optical galaxy and the position searched in the radio image.



Figure 3. The median 1.4GHz flux density detected within 0.75 arcseconds of the position of optical galaxies in the ACS HDF-N field, binned by z-band magnitude. The control sample incorporates a random 7 arcsecond shift.

Radio emission is clearly detected statistically for all z-band magnitude bins brighter than 26. Of the ∼2700 galaxies brighter than z=$24^{\rm mag}$, half (nearly 1400) will have radio flux densities of $\sim 4\mu$Jy or greater at 1.4GHz. A 4μJy source represents an $\sim 8\sigma$ detection for a future deep *e*-MERLIN+EVLA image; such galaxies which at present may only be investigated statistically will in the near future be imaged individually.

3.2. Radio Source Sizes

The angular sizes of the very weak radio sources can also be investigated statistically. In each magnitude bin the average radio source size can be derived from the radio flux densities found in annuli of width 0.25 arcseconds, over radii of 0.25 to 2 arcseconds centred on each galaxy position. These are shown in Figure 4. For detected systems brighter than z∼$25^{\rm mag}$, the average radio source radii are in the range 0.6 to 0.8 arcseconds implying that the next generation radio interferometers will need sub-arcsecond angular resolution. There is a marginal trend for the fainter radio sources to be slightly smaller. Radio source sizes may also be also be measured by fitting to the composite, stacked radio image in each magnitude bin. Similar sizes are derived by this latter method.



Figure 4. The average radio flux density contained within annuli of width 0.25 arcseconds over radii of 0.25 to 2 arcseconds centred at the position on each galaxy binned by z-band magnitude. For galaxies brighter than z$\sim$25$^{\rm mag}$, the average radio source radii are in the range 0.6 to 0.8 arcseconds.

3.3. Starburst Luminosities

Around 1000 of the $\sim$13000 galaxies in the 8.5 arcminute field have published spectroscopic redshifts. We have constructed the luminosity-redshift distribution for those starburst systems with measured redshifts. For this sub-set of starbursts, the detected radio flux density stacked by z-band magnitude does not differ significantly from the complete galaxy sample (at least for those bins brighter than 24$^{\rm mag}$); indicating that the sub-set of galaxies with measured redshifts do not appear to differ significantly from the complete galaxy set. We are thus able to derive the averaged luminosity distribution for those starburst galaxies brighter than 24$^{\rm mag}$

The very weak radio sources which are assumed to be dominated by starbursts have average properties which indicate that they are an extension of the weaker end of the distribution of starbursts detected as individual sources at higher radio flux densities. Their typical luminosities are still several times that of M82. However, it should be noted that these weak radio sources have been selected by optical z-band properties and will therefore exclude optically faint systems. Most of the individual high redshift sub-mm starburst systems imaged by Muxlow *et al.* (2005) were optically faint, and therefore will not be represented statistically in these results. Furthermore close galaxy pairs have also been excluded. Clearly individual imaging of these very faint systems will be required.

4. The Future

The new upgraded radio interferometers due to be commissioned in the next few years will be able to individually image these very faint systems. *e*-MERLIN, EVLA, and *e*-VLBI should be able to image in excess of 1000 individual starburst



Figure 5. Rest frame monochromatic luminosities for individual galaxies from Muxlow *et al.* (2005) and the statistical results presented here (filled diamonds). Sample cut-off levels of 40, 4, and 0.4μJy are shown. For comparison, two nearby starbursts (M82 & Arp220) are also shown together with HDF850.1, the brightest sub-mm *SCUBA* detection in the HDF-N

systems to flux densities of $\sim$4μJy at 1.4GHz with perhaps 150 to 200 objects at high redshift in a single field. For the even weaker radio source population, the new instruments will be able to extend this present statistical study to include many thousands of starburst systems with radio flux densities less then 1μJy. SKA and ALMA will ultimately extend this by an additional order of magnitude, With redshifts, improved spectral energy distribution templates and extinction-free star-formation rate indicators, it will become possible to solve for the cosmic star-formation history - the radio Madau diagram.

References

Condon, J. J., Condon, M. A., Gisler, G. & Puschell, J., 1982, ApJ, 252, 102
Garrett, M. A., *et al.*, 2001, A&A, 366, L5
Madau, P., Pozzetti, L., & Dickinson, M., 1998, ApJ, 498, 106
Muxlow, T. W. B., *et al.*, 2005 MNRAS, 358, 1159
Richards, E. A., Kellermann, K. I., Fomalont, E. B., Windhorst, R. A., Partridge, R. B., 1998, AJ, 116, 1039
Richards, E. A., 2000, ApJ, 533, 611
Seymour, N., McHardy, I.M., & Gunn, K.F., 2004, MNRAS, 352, 131
van der Kruit, P. C., 1973, A&A, 29, 263